

CENTRAL OHIO EAR, NOSE AND THROAT, INC.

**41 COMMERCE PARK DR.
WESTERVILLE, OH 43082**

**6860 PERIMETER DR. SUITE A
DUBLIN, OH 43016**

Three weeks (21 days) prior to your surgery date, DO NOT take any ASPIRIN, ANY OTHER MEDICATION CONTAINING ASPIRIN, or ANY ANTI-INFLAMMATORY MEDICATION. Aspirin and anti-inflammatory medications cause an abnormality of blood clotting. This effect may cause excessive bleeding at the time of surgery. TYLENOL may be used for pain. For any questions, contact your family doctor or call our office. Listed below are some drugs that are anti-inflammatory agents or contain aspirin.

**MEDICATIONS THAT SHOULD NOT BE TAKEN THREE WEEKS
PRIOR TO SURGERY AND TWO WEEKS AFTER SURGERY:**

**ALL ARTHRITIS & HEADACHE DRUGS
(ANTI-INFLAMMATORY)**

Advil
Ibuprofen
Medipren
Midol
Motrin
Nuprin
Rufen Tablets
Aleve (naproxyn)

ANALGESICS (PAIN TABLETS)

Alka-Seltzer effervescent
Anacin
Arthritis Pain Formula
Arthritis Strength Bufferin
Arthropan Liquid
A.S.A. compound
A.S.A. Enseals
Ascription A/D
Bayer Aspirin
Bayer Children's Aspirin
Buffaprin
Bufferin
Ecotrin
Empirin compound
Excedrin
Fiorinal
Goody's Headache Powder
Measurin
Phenaphen
Pyrroxate
Vanquish

MENSTRUAL PRODUCTS

Femcaps
Midol
Pamprin

Rev. 7/2008

COLD AND COUGH MEDICATIONS

Super Anahist
Alka Seltzer Plus
Bayer Decongestant
Bayer Children's Cold Tablets
Congespirin
Contact
Coricidin
Dristan
Sine-Aid
Sine-Off
St. Joseph cold tablets for children

SEDATIVES

Quiet World Tablets
Robaxisal
Zacrin
Trigesic
Tranquil
Cope
Dolcin
Dolor
Duragesic
Emagrin
Excedrin PM
Fizrin Powder
Nilain
Nytol Capsules & Tablets
Pabirin
Panodynes Analgesic
Percodan
Phensol

OTHER

**No alcohol 1 week prior and 2 weeks
after surgery**
Vitamin E
Clinoral
Dipridamole
Persantine/Coumadin
Ginseng